arXiv:1209.2520v1 [cond-mat.stat-mech] 12 Sep 2012

Second-Order Phase Transition in Heisenberg Model on Triangular Lattice with Competing Interactions

Ryo Tamura,[1,*] Shu Tanaka,[2,†] and Naoki Kawashima[3,‡]
[1] *International Center for Young Scientists, National Institute for Materials Science, 1-2-1, Sengen, Tsukuba-shi, Ibaraki, 305-0047, Japan*
[2] *Department of Chemistry, University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan*
[3] *Institute for Solid State Physics, University of Tokyo, 5-1-5, Kashiwanoha, Kashiwa-shi, Chiba, 227-8581, Japan*

(Dated: September 13, 2012)

We study the nature of phase transition in classical Heisenberg model on triangular lattice with competing interactions. In frustrated systems, various types of phase transitions occur depending on the lattice and spin symmetries. Recently, a first-order phase transition with threefold symmetry breaking in frustrated Heisenberg model on triangular lattice was discovered [R. Tamura and N. Kawashima, J. Phys. Soc. Jpn. **80**, 074008 (2011)]. To change the underlying symmetry, we introduce a uniaxial distortion effect into their model. We find the order of the phase transition changes into the second order when the spontaneous breaking of the twofold symmetry occurs. Moreover, dissociation of Z_2 vortex pairs occurs at the second-order phase transition point.

PACS numbers: 75.10.Hk, 64.60.De, 64.60.F-, 75.40.Mg

Geometrically frustrated magnets (GFMs) have been studied exhaustively in a wide range of areas[1–4]. In GFMs, occurrence of novel phase transitions associated with unconventional order parameters, existence of exotic ground states (GSs), and appearance of unconventional dynamical nature have been predicted theoretically and found experimentally. A number of examples in which theoretical calculation gives close agreement with experimental results of GFMs have been reported[5–9]. Moreover since recently there have been found new GFMs which exhibit theoretically uninvestigated nature[10, 11], theoretical studies of GFMs and investigations of resultant predictions have increased importance.

In geometrically frustrated systems, GS is not the conventional ferromagnetic state nor the Neél state in general. Non-collinear states such as the spiral-spin structure and the 120-degree structure become the GS. To describe such a spin structure, a characteristic degree of freedom such as chirality should be introduced. Moreover, a phase transition relating to the degree of freedom can take place in continuous spin systems even in two dimension where true long-range order of spins at finite temperature is prohibited by the Mermin-Wagner theorem[12]. For instance, the antiferromagnetic XY spin system on triangular lattice exhibits order-disorder transition associated with chirality[13], and the Z_2 vortex transition was reported in the antiferromagnetic Heisenberg spin system on triangular lattice[14]. Quest for peculiar phase transitions which never occur without frustration has been done for a long time.

It has been reported that some continuous spin systems with competing interactions exhibit a phase transition with spontaneous breaking of discrete symmetry corresponding to the lattice and spin symmetries[15–23]. In many cases, nature of phase transition is intuitively plausible from the symmetry that is broken at the transition point. However, a couple of unexpected phase transitions were reported. For example, in the J_1-J_2 XY model on square lattice, although a phase transition with twofold symmetry breaking occurs, it does not belong to the universality class of the two-dimensional Ising model[16]. Moreover, there has been recently found that the Heisenberg spin system on triangular lattice with the nearest-neighbor ferromagnetic interaction J_1 and the third nearest-neighbor antiferromagnetic interaction J_3 (J_1-J_3 model) exhibits a first-order phase transition with threefold symmetry breaking[18, 21]. This is also an unexpected phase transition since the order of the phase transition with breaking of the threefold symmetry in two dimension is usually the second order from the knowledge of the Potts model[24]. In this way, even now, geometrically frustrated systems have shown novel results.

In this paper, to investigate relationship between the symmetry that is broken at the transition point and nature of phase transition in continuous spin system with competing interactions, we introduce a uniaxial distortion parameter into the J_1-J_3 model[18, 21]. In the model, we find that a second-order phase transition with twofold symmetry breaking occurs at finite temperature for a finite range of the distortion parameter. Furthermore, we confirm the phase transition belongs to the universality class of the two-dimensional Ising model.

We consider the following Hamiltonian:

$$\begin{aligned}\mathcal{H}= \lambda J_1 \sum_{\langle i,j\rangle_{\mathrm{axis}\,1}} \mathbf{s}_i\cdot\mathbf{s}_j + J_1 \sum_{\langle i,j\rangle_{\mathrm{axis}\,2,3}} \mathbf{s}_i\cdot\mathbf{s}_j \\ + J_3 \sum_{\langle\langle i,j\rangle\rangle} \mathbf{s}_i\cdot\mathbf{s}_j, \qquad \lambda>0,\ J_3>0,\end{aligned} \tag{1}$$

where the first term represents nearest-neighbor interactions along the axis 1, the second term denotes nearest-neighbor interactions along the axes 2 and 3, and the summation in the third term takes over the third nearest-neighbor spin pairs (see Fig. 1(a)). The variable $\mathbf{s}_i$ is the three-dimensional vector spin of unit length, and λ is a uniaxial distortion parameter. Let the number of spins be $N(= L \times L)$, where L is the linear dimension. Hereafter we refer to this model as distorted J_1-J_3 model. In Ref. [21], the authors studied thermodynamic properties of the model for only $\lambda = 1$ and found that a first-order phase transition with breaking of the threefold symmetry occurs when the threefold symmetry is broken in the GS (*i.e.* $-4 < J_1/J_3 < 0$). Since in the GS for the other region of J_1/J_3 for $\lambda = 1$, no discrete symmetry is broken, we focus on the parameter region in this paper.

In general, the GS of the classical Heisenberg model can be represented by the wave vector $\mathbf{k}^*$ at which the Fourier transform of interactions $J(\mathbf{k})$ is minimized. In the distorted J_1-J_3 model, $J(\mathbf{k})$ is given by

$$\frac{J(\mathbf{k})}{NJ_3} = \frac{\lambda J_1}{J_3}\cos k_x + \frac{2J_1}{J_3}\cos\frac{k_x}{2}\cos\frac{\sqrt{3}k_y}{2} + \cos 2k_x + 2\cos k_x \cos\sqrt{3}k_y. \tag{2}$$

Notice that the spiral-spin state represented by $\mathbf{k}$ and that by $-\mathbf{k}$ are the same structure in the Heisenberg model. Figure 1(b) summarizes positions of $\mathbf{k}^*$ in the first Brillouin zone depending on λ. When $0 < \lambda < 1$, there are two wave vectors $\mathbf{k}^* = \pm(k_x^* \neq 0, k_y^* = 0)$. In this case, no discrete symmetry is broken in the GS. For only $\lambda = 1$, there are six wave vectors $\mathbf{k}^*$ and the threefold symmetry is broken in the GS as stated in Ref. [21]. If we increase λ from unity, there are four wave vectors $\mathbf{k}^*$. Then, the twofold symmetry is broken in the GS. Moreover for large λ ($> \lambda_0$ obtained by Eq. (2)), the wave vector $\mathbf{k}^*$ is given by $\mathbf{k}^* = \pm(k_x^* = 0, k_y^* \neq 0)$ and no discrete symmetry is broken in the GS. In this way, the symmetry that is broken in the GS can be changed by tuning λ. In the following, we focus on the parameter region $1 < \lambda < \lambda_0$, since nature of phase transition with breaking of the twofold symmetry has not been investigated in continuous spin system on triangular lattice. Note that the abovementioned all spin configurations have SO(3) symmetry reflecting the global rotation of all spins.

In order to obtain the equilibrium values of physical quantities with high accuracy, we perform the Monte Carlo simulation using the single-spin-flip heat-bath method and the over-relaxation method[25, 26]. First, we study the nature of phase transition in the distorted J_1-J_3 model with periodic boundary condition. To avoid extra stress caused by boundary effect, we use the parameter set such that the GS can be represented by the commensurate wave vector. We consider the case that the wave vectors $\mathbf{k}^*$ which minimize $J(\mathbf{k})$ are $\mathbf{k}^* = \pm(5\pi/18, 16\pi/18\sqrt{3})$ and $\pm(5\pi/18, -16\pi/18\sqrt{3})$. In the GS spin configuration in the case, the angle between nearest-neighbor spins along the axis 1 is 50-degree and that along one of the axes 2 and 3 is 55-degree and the other is 105-degree as shown in Fig. 1(a). We obtain $J_1/J_3 = -0.4926\cdots$ and $\lambda = 1.308\cdots$ using Eq. (2). Figure 1(c) shows temperature T dependence of the structure factor at $\mathbf{k}^* = (5\pi/18, 16\pi/18\sqrt{3})$ for $L = 144$ when the GS spin configuration is represented as the left panel of Fig. 1(a). The structure factor is defined as



FIG. 1. (Color online) (a) Schematic picture of two distinct GSs of the distorted J_1-J_3 model for $J_1/J_3 = -0.4926\cdots$ and $\lambda = 1.308\cdots$. The bold lines indicate λJ_1. (b) λ-dependence of $\mathbf{k}^*$ which are denoted by the circles in the first Brillouin zone depicted by the dotted hexagon. (c) Structure factor $S(\mathbf{k})$ at one of $\mathbf{k}^*$ as a function of temperature. Structure factors in the first Brillouin zone at several temperatures are also shown. The lattice size is $L = 144$.

$$S(\mathbf{k}) := \frac{1}{N}\sum_{i,j}\langle \mathbf{s}_i \cdot \mathbf{s}_j\rangle \, \mathrm{e}^{-i\mathbf{k}\cdot(\mathbf{r}_i - \mathbf{r}_j)}, \tag{3}$$

where the summation takes over all of the spin pairs and $\mathbf{r}_i$ represents the position vector of the i-th site. Throughout the paper, $\langle \mathcal{O} \rangle$ denotes the equilibrium value of the physical quantity $\mathcal{O}$ and the Boltzmann constant is set to unity. As temperature decreases, $S(\mathbf{k}^*)$ monotonically increases. The structure factor in the first Brillouin zone at several temperatures is also shown in Fig. 1(c), which suggests the twofold symmetry is broken at low temperature[27].

We calculate temperature dependence of specific heat for $L = 144 - 288$. The specific heat is calculated as

$$C = (\langle E^2\rangle - \langle E\rangle^2)/T^2, \tag{4}$$

where E is the internal energy per site. Figure 2(a) shows the specific heat as a function of temperature and indicates the existence of a phase transition at finite temperature. To confirm the occurrence of a spontaneous twofold symmetry breaking, we define an order parameter m by using a local quantity $\kappa^{(t)}$ at each upward elementary triangle t depicted by the shaded triangles in Fig. 1(a):

$$\kappa^{(t)} := \mathbf{s}_1^{(t)} \cdot (\mathbf{s}_2^{(t)} - \mathbf{s}_3^{(t)}), \qquad m := \sum_t \kappa^{(t)}/N, \tag{5}$$

where the definition of the subscript of spins is shown in Fig. 2(b) and the summation takes over all upward elementary triangles. Since the order parameter represents the difference between the relative angle of nearest-neighbor spins along the axis 2 and that along the axis 3, it can appropriately describe the twofold symmetry breaking in our model. Figures 2(b) and (c) show the temperature dependence of $\langle m^2\rangle$ and $U_4 := \langle m^4\rangle/\langle m^2\rangle^2$, respectively. These figures indicate the existence of the second-order phase transition with breaking of the twofold symmetry at $T_c/J_3 = 0.4950(5)$ where the Binder ratios for some lattice sizes cross.

Since the spiral-spin states in the GS of antiferromagnetic Heisenberg spin systems on triangular lattice have SO(3) symmetry, the order parameter space is isomorphic to P_3 and the point defect is $\pi_1(\mathrm{P}_3) = Z_2$. Because of the existence of Z_2 point defect, the dissociation of Z_2 vortex pairs occurs at finite temperature[14, 28]. The Z_2 vortex labeled by 0 or 1 is defined by the vortex configuration of spin and vector chirality. In order to confirm the dissociation of Z_2 vortex pairs in our model, we calculate the number density of Z_2 vortices n_v by using the same manner as in Ref. [14]. A plot of $\ln n_\mathrm{v}$ versus J_3/T in our model is shown in Fig. 2(d), and it is confirmed that n_v obeys well the Arrhenius law below T_c. This result indicates that the dissociation of Z_2 vortex pairs occurs at the second-order phase transition point. Note that Z_2 vortex pairs dissociate at the phase transition point was also found for $\lambda = 1$[21], but, in the case, the order of the phase transition is the first order. Hence, it is concluded that the phase transition associated with breaking of discrete symmetry and the dissociation of Z_2 vortex pairs occur simultaneously regardless of the order of the phase transition in our model.

To clarify the universality class of the phase transition, we perform the finite-size scaling using the following relations:

$$U_4 \propto f((T - T_\mathrm{c})L^{1/\nu}),\ \chi \propto L^{2-\eta} g((T - T_\mathrm{c})L^{1/\nu}), \tag{6}$$

where the susceptibility χ is defined as $\chi := NJ_3\langle m^2\rangle/T$ and $f(\cdot)$ and $g(\cdot)$ are scaling functions. The obtained results using $\nu = 1$ and $\eta = 1/4$ which are the critical exponents of the two-dimensional Ising model and the obtained T_c are shown in Figs. 2(e) and (f). Since all the data collapse onto scaling functions, it is confirmed that the second-order phase transition in our model belongs to the universality class of the Ising model.



FIG. 2. (Color online) Temperature dependence of equilibrium physical quantities of the distorted J_1-J_3 model for $J_1/J_3 = -0.4926\cdots$ and $\lambda = 1.308\cdots$. (a) Specific heat C. (b) Square of the order parameter $\langle m^2\rangle$. (c) Binder ratio U_4. (d) Log of number density of Z_2 vortex n_v versus J_3/T. The dotted vertical line indicates the transition temperature $T_\mathrm{c}/J_3 = 0.4950(5)$. (e) and (f) Finite-size scaling of the Binder ratio U_4 and that of the susceptibility χ using the critical exponents of the two-dimensional Ising model ($\nu = 1$ and $\eta = 1/4$) and the transition temperature. Error bars are omitted for clarity since their sizes are smaller than the symbol sizes.

Next, to obtain the relationship between λ and T_c, we consider the case of $J_1/J_3 = -0.7342\cdots$ which is used in Ref. [21] with changing the value of λ. For $\lambda = 1$, the model exhibits a first-order phase transition with breaking of the threefold symmetry at $T_\mathrm{c}/J_3 = 0.4746(1)$[21]. We here study the nature of phase transition of the distorted J_1-J_3 model with open boundary condition. From the analysis of the GS as explained before, a phase transition with breaking of the twofold symmetry is expected to take place for $1 < \lambda < \lambda_0 (= 2.8155\cdots)$ in this case. By analyzing the Binder ratio, we obtain λ-dependence of transition temperatures as depicted in Fig. 3(a). An enlarged view near $\lambda = 1$ is shown in the inset of Fig. 3(a). This figure indicates the transition temperature near $\lambda = 1$ smoothly connects to the transition temperature for $\lambda = 1$ and the transition temperature goes continuously to zero when $\lambda \to \lambda_0$. Figures 3(b) and (c) represent the finite-size scaling analysis of the Binder ratio and that of the susceptibility for $\lambda = 1.5$ using $\nu = 1$, $\eta = 1/4$, and $T_\mathrm{c}/J_3 = 0.5521(1)$ as well as



FIG. 3. (Color online) (a) Phase diagram of the distorted J_1-J_3 model for $J_1/J_3 = -0.7342\cdots$. The inset is an enlarged view. The open square indicates the transition temperature for $\lambda = 1$ where a first-order phase transition with threefold symmetry breaking occurs[21]. The solid circles represent transition temperatures at which a second-order phase transition with twofold symmetry breaking occurs. (b) and (c) Finite-size scaling of the Binder ratio U_4 and that of the susceptibility χ for $\lambda = 1.5$ using $\nu = 1$ and $\eta = 1/4$ which are the critical exponents of the two-dimensional Ising model. The transition temperature is decided as $T_c/J_3 = 0.5521(1)$. Error bars are omitted for clarity since their sizes are smaller than the symbol sizes.

the previous case. In this case, all the data collapse onto scaling functions. By the same procedure as the previous case, we obtain transition temperatures which are indicated by the solid circles in Fig. 3(a). Thus, we conclude that a second-order phase transition with breaking of the twofold symmetry occurs and it belongs to the two-dimensional Ising model universality class in the parameter region where twofold symmetry is broken in the GS within the calculated parameters.

In this paper, we investigated nature of phase transition in classical Heisenberg model on triangular lattice with three types of interactions: (i) the uniaxially distorted nearest-neighbor ferromagnetic interaction along the axis 1 (λJ_1), (ii) the nearest-neighbor ferromagnetic interaction along the axes 2 and 3 (J_1), and (iii) the third nearest-neighbor antiferromagnetic interaction (J_3). In Ref. [21], the authors studied for only $\lambda = 1$ (no distortion) and they found that a first-order phase transition with threefold symmetry breaking occurs when the ground state is a spiral-spin state with breaking of the threefold symmetry. To investigate a uniaxial distortion effect for phase transition, properties of the ground state were studied and we found that the order of the phase transition changes into the second order when the twofold symmetry is broken in the ground state. The universality class of the second-order phase transition is the same as that of the two-dimensional Ising model. We also confirmed that the dissociation of Z_2 vortex pairs occurs at the second-order phase transition temperature. Although a second-order phase transition occurs in considered λ, there is still an open problem whether a first-order phase transition with breaking of the twofold symmetry occurs at very close to $\lambda = 1$. This is because the property of phase transition is decided by not only the symmetry that is broken at the transition point but also the structure of density of states[29]. If a first-order phase transition with breaking of the twofold symmetry occurs, a tricritical point should exist and to study its properties such as universality class becomes an important topic. From our observation, it is difficult to decide the order of the phase transition near $\lambda = 1$ since the size dependence of physical quantities are significant and we should calculate very large systems with high accuracy.

Our model exhibits both a first-order phase transition and a second-order phase transition by tuning the uniaxial distortion parameter λ. We believe that the obtained nature is not restricted to the considered model. This is because a first-order phase transition accompanied with threefold symmetry breaking was also shown in geometrically frustrated continuous spin systems *e.g.* the J_1-J_2 Heisenberg model on triangular lattice and that on honeycomb lattice[20, 22]. Moreover, the uniaxial distortion parameter λ can be experimentally changed by pressure or/and chemical substitution. Then such nature is expected to be realized in many geometrically frustrated magnets.

The authors are partially supported by National Institute for Materials Science (NIMS), Grand-in-Aid for JSPS Fellows (23-7601) and Scientific Research (B) (22340111), the Computational Materials Science Initiative (CMSI). Numerical calculations were performed on supercomputers at the Institute for Solid State Physics, University of Tokyo.

---

[*] tamura.ryo@nims.go.jp
[†] shu-t@chem.s.u-tokyo.ac.jp
[‡] kawashima@issp.u-tokyo.ac.jp

[1] G. Toulouse, Commun. Phys. (London) **2**, 115 (1977).

[2] R. Liebmann, *Statistical Mechanics of Periodic Frustrated Ising Systems* (Springer-Verlag, Berlin/Heidelberg GmbH, Heidelberg, 1986).

[3] H. Kawamura, J. Phys.: Condens. Matter **10**, 4707 (1998).

[4] *Frustrated Spin Systems*, edited by H. T. Diep (World Scientific, Singapore, 2005).

[5] H. Kageyama, K. Yoshimura, R. Stern, N. V. Mushnikov, K. Onizuka, M. Kato, K. Kosuge, C. P. Slichter, T. Goto, and Y. Ueda, Phys. Rev. Lett. **82**, 3168 (1999).

[6] A. P. Ramirez, A. Hayashi, R. J. Cava, R. Siddharthan, and B. S. Shastry, Nature **399**, 333 (1999).

[7] B. C. den Hertog and M. J. P. Gingras, Phys. Rev. Lett. **84**, 3430 (2000).

[8] C. Castelnovo, R. Moessner, and S. L. Sondhi, Nature **451**, 42 (2008).

[9] R. Ishii, S. Tanaka, K. Onuma, Y. Nambu, M. Tokunaga, T. Sakakibara, N. Kawashima, Y. Maeno, C. Broholm, D. P. Gautreaux, J. Y. Chan, and S. Nakatsuji, Euro-

phys. Lett. **94**, 17001 (2011).

[10] H. Yoshida, J. Yamaura, M. Isobe, Y. Okamoto, G. J. Nilsen, and Z. Hiroi, Nature Communications **3**, 860 (2012).

[11] S. Nakatsuji, K. Kuga, K. Kimura, R. Satake, N. Katayama, E. Nishibori, H. Sawa, R. Ishii, M. Hagiwara, F. Bridges, T. U. Ito, W. Higemoto, Y. Karaki, M. Halim, A. A. Nugroho, J. A. Rodriguez-Rivera, M. A. Green, and C. Broholm, Science **336**, 559 (2012).

[12] N. D. Mermin and H. Wagner, Phys. Rev. Lett. **17**, 1133 (1966).

[13] S. Miyashita and H. Shiba, J. Phys. Soc. Jpn. **53**, 1145 (1984).

[14] H. Kawamura and S. Miyashita, J. Phys. Soc. Jpn. **53**, 4138 (1984).

[15] P. Chandra, P. Coleman, and A. I. Larkin, Phys. Rev. Lett. **64**, 88 (1990).

[16] D. Loison and P. Simon, Phys. Rev. B **61**, 6114 (2000).

[17] C. Weber, L. Capriotti, G. Misguich, F. Becca, M. Elhajal, and F. Mila, Phys. Rev. Lett. **91**, 177202 (2003).

[18] R. Tamura and N. Kawashima, J. Phys. Soc. Jpn. **77**, 103002 (2008).

[19] E. M. Stoudenmire, S. Trebst, and L. Balents, Phys. Rev. B **79**, 214436 (2009).

[20] S. Okumura, H. Kawamura, T. Okubo, and Y. Motome, J. Phys. Soc. Jpn. **79**, 114705 (2010).

[21] R. Tamura and N. Kawashima, J. Phys. Soc. Jpn. **80**, 074008 (2011).

[22] T. Okubo, S. Chung, and H. Kawamura, Phys. Rev. Lett. **108**, 017206 (2012).

[23] S. Jin, A. Sen, and A. W. Sandvik, Phys. Rev. Lett. **108**, 045702 (2012).

[24] F. Y. Wu, Rev. Mod. Phys. **54**, 235 (1982).

[25] M. Creutz, Phys. Rev. D **36**, 515 (1987).

[26] K. Kanki, D. Loison, and K. D. Schotte, Eur. Phys. J. B **44**, 309 (2005).

[27] Note that, since the long-range order of spins at finite-temperature is prohibited by the Mermin-Wagner theorem in our model, $S(\mathbf{k})$ for arbitrary $\mathbf{k}$ should be vanished in the thermodynamic limit. Then, $S(\mathbf{k})$ as shown in Fig. 1(c) is just an indication for finite size systems. In order to study the phase transition with discrete symmetry breaking at finite temperature in our model, we should define an appropiate order parameter instead of $S(\mathbf{k})$ and investigate its thermodynamic properties.

[28] H. Kawamura, A. Yamamoto, and T. Okubo, J. Phys. Soc. Jpn. **79**, 023701 (2010).

[29] R. Tamura, S. Tanaka, and N. Kawashima, Prog. Theor. Phys. **124**, 381 (2010).